

Discontinuous Galerkin method for an N-dimensional spherically symmetric Poisson equation

Yuki Chiba

Graduate School of Mathematical Sciences, University of Tokyo

Joint work with Norikazu Saito

International Workshop on Numerical Methods for Partial Differential
Equations

The Hong Kong Polytechnic University

March 26, 2018

Contents

1. Introduction
2. DG Scheme
3. Analysis of Scheme
4. Numerical Results
5. Modified DG Scheme

1 . Introduction

Introduction

In the theory of PDE, it is known several critical phenomena with critical value related to space dimension N .

Example: Positive solution of semilinear elliptic equation

$N \geq 3$, $\Omega \subset \mathbb{R}^N$: smooth bounded domain.

$$\begin{cases} \Delta u + |u|^{p-1} u = 0 & (x \in \Omega) \\ u = 0 & (x \in \partial\Omega) \end{cases}$$

If $1 < p < p_s = (N + 2)/(N - 2)$, positive solution exists. If $p > p_s$, positive solution may not exist.

If we can make numerical method for nonlinear PDEs in higher dimension, it is useful for studying critical phenomena through experimental consideration.

Model Problem

Consider the Poisson equation in N -dimensional ball

$$B_R = \{\xi \in \mathbb{R}^N \mid |\xi|_{\mathbb{R}^N} < R\}.$$

Poisson equation

$$\begin{cases} -\Delta_{\xi}U(\xi) + Q(\xi)U(\xi) = F(\xi) & (\xi \in B_R) \\ U(\xi) = 0 & (\xi \in \partial B_R) \end{cases} \quad (1)$$

Set $x = |\xi|$ and we assume that coefficient function Q and F are spherically symmetric $Q(\xi) = \hat{q}(x)$, $F(\xi) = \hat{f}(x)$.

Then, (1) is reduced to next equation.

$$\begin{cases} -\frac{1}{x^{N-1}} (x^{N-1}\hat{u}_x)_x + \hat{q}\hat{u} = \hat{f} & (x \in I = (0, R)) \\ \hat{u}_x(0) = \hat{u}(R) = 0 \end{cases} \quad (2)$$

Model Problem

In previous study, there are two FEMs using weight function to eliminate singularity. (cf. K, Ericsson and V, Thomée. 1984)

1. Using weight function x^{N-1}

$$-(x^{N-1}\hat{u}_x)_x + x^{N-1}\hat{q}\hat{u} = x^{N-1}\hat{f}$$

2. Using weight function x

$$-(x\hat{u}_x)_x + (2 - N)\hat{u}_x + x\hat{q}\hat{u} = x\hat{f}$$

In this study, we apply Discontinuous Galerkin (DG) method to second case.

2 . DG Scheme

Generally, consider the following problem.

Diffusion-convection equation

$$\begin{cases} -(\nu u_x)_x + bu_x + qu = f & (x \in I) \\ u_x(0) = u(R) = 0 \end{cases} \quad (3)$$

$$\nu(x) = x, \quad b \leq 0: \text{Const.}, \quad q, f \in L^2(I), \quad q(x) > 0 \quad (x \in I)$$

Division $\mathcal{T}_h = \{K_i\}_{i \in \Lambda}$ of I is defined by below.

$$0 = x_1 < x_2 < \cdots < x_i < \cdots < x_n = R$$

$$K_i = (x_i, x_{i+1}), \quad h_i = |K_i| = x_{i+1} - x_i,$$

$$h = \max_{i \in \Lambda} h_i, \quad \Lambda = \{1, 2, \dots, n-1\}$$

$$e_i = \min\{h_i, h_{i-1}\} \quad (i = 2, \dots, n-1), \quad e_n = h_{n-1}$$

Function space

$$H^m(\mathcal{T}_h) = \{v \in L^2(I) \mid v|_{K_i} \in H^m(K_i) \ (i \in \Lambda)\}$$

$$V_h = V_h^k = \{v \in L^2(I) \mid v|_{K_i} \in \mathcal{P}^k(K_i) \ (i \in \Lambda)\}$$

Notation: For $v \in H^1(\mathcal{T}_h)$, $v^i = v|_{K_i}$ ($i \in \Lambda$)

$$\nu_i = \nu(x_i) = x_i, \quad (u, v)_i = \int_{x_i}^{x_{i+1}} uv \, dx$$

$$[[v]]_i = \begin{cases} -v^1(x_1) & (i = 1) \\ v^{i-1}(x_i) - v^i(x_i) & (2 \leq i \leq n-1) \\ v^{n-1}(x_n) & (i = n) \end{cases}$$

$$\langle\langle v \rangle\rangle_i = \begin{cases} v^1(x_1) & (i = 1) \\ \frac{v^{i-1}(x_i) + v^i(x_i)}{2} & (2 \leq i \leq n-1) \\ v^{n-1}(x_n) & (i = n) \end{cases}$$

DG Scheme

$$\begin{aligned} \text{Find } u_h \in V_h \quad \text{s.t.} \\ a_h(u_h, v) = a_h^d(u_h, v) + a_h^{\text{cr}}(u_h, v) = (f, v) \quad (\forall v \in V_h) \end{aligned} \quad (4)$$

$$\begin{aligned} a_h^d(u, v) &= \sum_{i=1}^{n-1} (\nu u_x, v_x)_i - \sum_{i=2}^n \nu_i \langle\langle u_x \rangle\rangle_i [[v]]_i \\ &\quad - \alpha \sum_{i=2}^n \nu_i \langle\langle v_x \rangle\rangle_i [[u]]_i + \sum_{i=2}^n \frac{\nu_i \sigma}{e_i} [[u]]_i [[v]]_i \\ a_h^{\text{cr}}(u, v) &= - \sum_{i=1}^{n-1} (bu, v_x)_i + \sum_{i=1}^{n-1} b \langle\langle u \rangle\rangle_i [[v]]_i + \sum_{i=2}^{n-1} \frac{1}{2} |b| [[u]]_i [[v]]_i + \sum_{i=1}^{n-1} (qu, v)_i \\ (f, v) &= \sum_{i=1}^{n-1} (f, v)_i \end{aligned}$$

3 . Analysis of Scheme

We introduce following DG norms.

$$\|v\|_d^2 = \sum_{i=1}^{n-1} (\nu v_x, v_x)_i + \sum_{i=2}^n \frac{\nu_i \sigma}{e_i} \llbracket v \rrbracket_i^2, \quad \|v\|_{d,*}^2 = \|v\|_d^2 + \sum_{i=1}^{n-1} h_i^2 (\nu v_{xx}, v_{xx})_i$$

$$\|v\|_{cr}^2 = \sum_{i=1}^{n-1} (qv, v)_i + \sum_{i=1}^n \frac{1}{2} |b| \llbracket v \rrbracket_i^2, \quad \|v\|_{cr,*}^2 = \|v\|_{cr}^2 + \sum_{i=1}^{n-1} |b| \langle\langle v \rangle\rangle_i^2$$

$$\| \|v\| \|^2 = \|v\|_d^2 + \|v\|_{cr}^2, \quad \| \|v\| \|^2_* = \|v\|_{d,*}^2 + \|v\|_{cr,*}^2$$

Assume that division $\{\mathcal{T}_h\}_h$ is quasi-uniform.

$$\exists \theta_0 > 0 \quad \text{s.t.} \quad 0 < \frac{h_i}{h_j} \leq \theta_0 \quad (1 \leq \forall i, j \leq n, \quad \forall \mathcal{T}_h \in \{\mathcal{T}_h\}_h) \quad (\text{A1})$$

Lem 1 (Trace inequality)

There exists a positive constant $C_1 = C_1(\theta_0)$ satisfying

$$\nu_i(v_x^i(x_i))^2 \leq C_1(h_i^{-1}(\nu v_x, v_x)_i + h_i(\nu v_{xx}, v_{xx})_i)$$

$$\nu_i(v_x^i(x_{i+1}))^2 \leq C_1(h_i^{-1}(\nu v_x, v_x)_i + h_i(\nu v_{xx}, v_{xx})_i).$$

Lem 2 (Inverse inequality)

Let $K = (s, t) \subset \mathbb{R}_{>0}$ be interval and set $\rho = t - s > 0$. Then, There exists a positive constant $C_2 = C_2(k)$ satisfying

$$\int_K x v_{xx}^2 dx \leq C_2 \rho^{-2} \int_K x v_x^2 dx \quad (v \in \mathcal{P}^k).$$

Lem 3 (Continuity and coercivity of a_h^d)

(i) For all $\alpha \in \mathbb{R}$ and $\sigma > 0$, there exists a positive constant $C_d > 0$ independent of h satisfying

$$a_h^d(u, v) \leq C_d \|u\|_{d,*} \|v\|_d \quad (u \in H^2(\mathcal{T}_h), v \in V_h).$$

(ii) There exists a positive constant $\sigma_* > 0$ independent of h satisfying below. If $\sigma \geq \sigma_*$, then

$$a_h^d(v, v) \geq \frac{1}{2} \|v\|_d^2 \quad (v \in V_h).$$

Proof

(i) By Cauchy-Schwarz's inequality,

$$\begin{aligned}
 |a_h^d(u, v)| &\leq \left(\sum_{i=1}^{n-1} (\nu u_x, u_x)_i^2 + \sum_{i=2}^n \frac{\nu_i e_i}{\sigma} \langle\langle u_x \rangle\rangle_i^2 + (1 + |\alpha|) \sum_{i=2}^n \frac{\nu_i \sigma}{e_i} [u]_i^2 \right)^{1/2} \\
 &\quad \cdot \left(\sum_{i=1}^{n-1} (\nu v_x, v_x)_i^2 + \sum_{i=2}^n \frac{\nu_i e_i}{\sigma} \langle\langle v_x \rangle\rangle_i^2 + (1 + |\alpha|) \sum_{i=2}^n \frac{\nu_i \sigma}{e_i} [v]_i^2 \right)^{1/2}.
 \end{aligned}$$

Using trace inequality and inverse inequality, we get the estimate.

Proof

(ii) Using Cauchy-Schwarz's inequality, trace inequality and inverse inequality, for $\delta > 0$,

$$a_h^d(v, v) \geq \left(1 - \frac{1 + |\alpha|}{2\delta\sigma} C_1(1 + C_2)\right) \sum_{i=1}^{n-1} (\nu v_x, v_x)_i \\ + \left(1 - \frac{1 + |\alpha|}{2}\delta\right) \sum_{i=2}^n \frac{\nu_i \sigma}{e_i} \llbracket v \rrbracket_i^2.$$

Choosing $\delta = 1/(1 + |\alpha|)$ and $\sigma \geq \sigma_* = C_1(1 + C_2)(1 + |\alpha|)/\delta$, we get the estimate.

Analysis of Convection Term a_h^{cr}

For each $i \in \Lambda$, we define $P_{K_i} : L^1(K_i) \rightarrow \mathcal{P}^k(K_i)$ as locally L^2 projection operator.

We define globally L^2 projection operator $P_h : L^1(I) \rightarrow V_h$ by $(P_h v)|_{K_i} = P_{K_i} v$ ($i \in \Lambda$).

Lem 4 (Continuity and covercivity of a_h^{cr})

(i) There exists a positive constant $C_{\text{cr}} > 0$ independent of h satisfying

$$a_h^{\text{cr}}(u - P_h u, v) \leq C_{\text{cr}} \|u - P_h u\|_{\text{cr},*} \|v\|_{\text{cr}} \quad (u \in H^2(\mathcal{T}_h), v \in V_h).$$

(ii) Following inequality holds.

$$a_h^{\text{cr}}(v, v) \geq \frac{1}{2} \|v\|_{\text{cr}}^2 \quad (v \in V_h)$$

Proof

(i) Set $\phi = u - P_h u$. Since $v_x \in \mathcal{P}^{k-1}$, $(\phi, v_x)_i = 0$. Using Cauchy-Schwarz's inequality,

$$\begin{aligned}
 a_h^{\text{cr}}(\phi, v) &\leq \left(\sum_{i=1}^{n-1} |b| \langle\langle \phi \rangle\rangle_i^2 + \sum_{i=2}^{n-1} \frac{1}{2} |b| [\![\phi]\!]_i^2 + \sum_{i=1}^{n-1} (q\phi, \phi)_i \right)^{1/2} \\
 &\quad \cdot \left(\sum_{i=1}^{n-1} |b| [\![v]\!]_i^2 + \sum_{i=2}^{n-1} \frac{1}{2} |b| [v]_i^2 + \sum_{i=1}^{n-1} (qv, v)_i \right)^{1/2} \\
 &\leq 3 \|\phi\|_{\text{cr},*} \|v\|_{\text{cr}}.
 \end{aligned}$$

Proof

(ii) Using integration by parts,

$$\begin{aligned} - \sum_{i=1}^{n-1} (bv, v_x)_i &= \sum_{i=1}^{n-1} [(bv_x, v)_i - b(v^i(x_{i+1}))^2 - v^i(x_i)^2] \\ &= \sum_{i=1}^{n-1} (bv_x, v)_i + bv^1(x_1)^2 - bv^{n-1}(x_n)^2 - 2 \sum_{i=2}^{n-1} b \langle\langle u \rangle\rangle_i [v]_i. \end{aligned}$$

So,

$$- \sum_{i=1}^{n-1} (bv, v_x)_i = [bv^1(x_1)^2 - bv^{n-1}(x_n)^2] / 2 - \sum_{i=2}^{n-1} b \langle\langle u \rangle\rangle_i [v]_i.$$

Substituting this for $a_h^{\text{cr}}(v, v)$, we get the estimate.

Lem 5 (Continuity and Coercivity of a_h)

(i) For all $\alpha \in \mathbb{R}$ and $\sigma > 0$, there exists a positive constant $C_{\text{dcr}} > 0$ independent of h satisfying

$$a_h(u - P_h u, v) \leq C_{\text{dcr}} \| \|u - P_h u\|_* \| \|v\| \| \quad (u \in H^2(\mathcal{T}_h), v \in V_h).$$

(ii) There exists a positive constant σ_* independent of h satisfying below. If $\sigma \geq \sigma_*$, then

$$a_h(v, v) \geq \frac{1}{2} \| \|v\| \|^2 \quad (v \in V_h).$$

Thm 1

Let $u \in H^2(I)$ be the solution of (3). Assume that $\sigma \geq \sigma_*$. Then, there exists a unique solution $u_h \in V_h$ of DG Scheme (4), and it satisfies Galerkin orthogonality

$$a_h(u - u_h, v) = 0 \quad (v \in V_h).$$

In addition, the following estimate holds.

$$\| \| u - u_h \| \| \leq (1 + 2C_{\text{dcr}}) \| \| u - P_h u \| \|_* . \quad (5)$$


If u is sufficiently smooth, the order of error is $O(h^k)$.

4 . Numerical Results


Numerical Results

$I = (0, 1)$, $N = 3, 100$, $b = 2 - N$, $\alpha = 1$, $\sigma = 20$

(i) $u(x) = \cos \frac{\pi}{2}x$



Error of $\| \cdot \|$ using \mathcal{P}^1 ($k = 1$)




Error of $\| \cdot \|$ using \mathcal{P}^2 ($k = 2$)

The order of error is $O(h^k)$ for sufficiently small h .


Numerical Results

$I = (0, 1)$, $N = 3, 100$, $b = 2 - N$, $\alpha = 1$, $\sigma = 20$

(ii) $u(x) = x^{7/4} - 1$



Error of $\| \cdot \|$ using \mathcal{P}^1 ($k = 1$)




Error of $\| \cdot \|$ using \mathcal{P}^2 ($k = 2$)

The order of error is $O(h)$ if $k = 1$ and $O(h^{7/4})$ if $k = 2$.


Observations of Other Norms

$I = (0, 1)$, $N = 3, 100$, $b = 2 - N$, $\alpha = 1$, $\sigma = 20$, \mathcal{P}^1 element

(i) $u(x) = \cos \frac{\pi}{2}x$



Error for $N = 3$




Error for $N = 100$


$\| \cdot \|$ norm is $O(h)$, the others are $O(h^2)$.

Observations of Other Norms

$I = (0, 1)$, $N = 3, 100$, $b = 2 - N$, $\alpha = 1$, $\sigma = 20$, \mathcal{P}^1 element
(ii) $u(x) = x^{7/4} - 1$



Error for $N = 3$




Error for $N = 100$


L^2 and weighted L^2 norm are $O(h^2)$, $\|\cdot\|$ norm is $O(h)$, L^∞ norm is $O(h^{7/4})$.

Observations of Other Norms

$I = (0, 1)$, $N = 3, 100$, $b = 2 - N$, $\alpha = 1$, $\sigma = 20$, \mathcal{P}^1 element
(iii) $u(x) = x^{5/4} - 1$



Error for $N = 3$




Error for $N = 100$


L^2 norm is $O(h^{7/4})$, weighted L^2 norm is $O(h^2)$, $\|\cdot\|$ norm is $O(h)$, L^∞ norm is $O(h^{5/4})$.

Observations of Other Norms

$I = (0, 1)$, $N = 3, 100$, $b = 2 - N$, $\alpha = 1$, $\sigma = 20$, \mathcal{P}^1 element
(iv) $u(x) = x^{3/4} - 1$



Error for $N = 3$



Error for $N = 100$

L^2 norm is $O(h^{5/4})$, weighted L^2 norm is $O(h^{7/4})$, $\|\cdot\|$ norm and L^∞ norm are $O(h^{3/4})$.

5 . Modified DG Scheme

Modified DG Scheme

$$\begin{aligned} &\text{Find } u_h \in V_h \quad \text{s.t.} \\ &b_h(u_h, v) = (f, v) \quad (\forall v \in V_h) \end{aligned} \tag{6}$$

$$\begin{aligned} b_h(u, v) = & \sum_{i=1}^{n-1} (\nu u_x, v_x)_i - \sum_{i=2}^n \nu_i \langle\langle u_x \rangle\rangle_i \llbracket v \rrbracket_i + \sum_{i=2}^n \frac{\nu_i \sigma}{e_i} \llbracket u \rrbracket_i \llbracket v \rrbracket_i \\ & + \sum_{i=1}^{n-1} (b u_x, v)_i - \sum_{i=2}^n b \llbracket u \rrbracket_i \langle\langle v \rangle\rangle_i + \sum_{i=2}^{n-1} \frac{1}{2} |b| \llbracket u \rrbracket_i \llbracket v \rrbracket_i + \sum_{i=1}^{n-1} (q u, v)_i \end{aligned}$$

Thm 2

Let $u \in H^2(I)$ be the solution (3). Assume that $\sigma \geq \sigma_*$.

Then, there exists a unique solution $u_h \in V_h$ of DG Scheme (6), and it satisfies Galerkin orthogonality.

In addition, there exists a positive constant $C > 0$ independent of h satisfying

$$\|u - u_h\|_{L^\infty(I)} \leq C(h \inf_{\chi \in V_h} \|u - \chi\|_{\text{DG}, \infty, 0} + \max_{i \in \Lambda} |(u - u_h)^i(x_{i+1})|)$$

$$\max_{i \in \Lambda} \|(u - u_h)_x\|_{L^\infty(K_i)} \leq C(\inf_{\chi \in V_h} \|u - \chi\|_{\text{DG}, \infty, 0} + \max_{i \in \Lambda} |(u - u_h)^i(x_{i+1})|)$$


for sufficiently small h . Moreover, if $q = 0$, we have

$$\max_{i \in \Lambda} \|(u - u_h)_x\|_{L^\infty(K_i)} \leq C \inf_{\chi \in V_h} \|u - \chi\|_{\text{DG}, \infty, 0}.$$


Modified DG Scheme - Numerical Result-

$I = (0, 1)$, $N = 100$, $b = 2 - N$, $\sigma = 20$

$$u(x) = \cos \frac{\pi}{2}x$$



Error using \mathcal{P}^1 ($k = 1$)



Error using \mathcal{P}^2 ($k = 2$)

L^∞ norm is $O(h^{k+1})$ and piecewise $W^{1,\infty}$ semi norm is $O(h^k)$.

Conclusion

- We have introduced DG schemes for a singular-perturbation elliptic problem derived from a spherical symmetric Poisson equation in the N dimensional ball. We have derived error estimates in the DG energy norm.
- We have confirmed the rate of convergence by numerical experiments. Optimal orders (depending on the regularity of solutions) were actually observed.
- Some point-wise estimates were obtained for a modified DG scheme.
- In the future work, I will apply the results to evolution equations and extend to nonlinear problems.